

Supplementary table 1: Risk factors, clinical status and treatment between participants from person-to-person and telephone survey

Prevalence n (%)	Person-to-person Survey N: 666	Telephone Survey N: 430	p value
Risk factors			
Smoker	92 (13.8)	45 (10.5)	0.111
Hypercholesterolemia	554 (84.1)	305 (71.9)	<0.001
Arterial hypertension	585 (88.8)	348 (82.3)	0.002
Diabetes mellitus	216 (32.8)	143 (33.6)	0.779
Clinical status			
Angina pectoris	149 (22.4)	67 (15.8)	0,008
Dyspnoea (CF II-IV)	194 (29.2)	142 (64.5)	<0,001
Treatment			
Betablockers	503 (75,6)	214 (49,8)	<0,001
Statins	575 (86,5)	234 (54,5)	<0,001
Antiplatelets	586 (88,3)	297 (69,2)	<0,001
Anticoagulants	55 (8,3)	56 (13,1)	0,011
ACE inh	263 (39,7)	122 (28,4)	<0,001
ARB	188 (28,4)	62 (14,5)	<0,001
Diuretics	251 (37,9)	133 (31,1)	0,022
Calcium antagonists	139 (21,0)	58 (13,6)	0,002
NTG transdermal	75 (11,3)	55 (12,9)	0,455
Oral antidiabetics	54 (8,2)	26 (6,1)	0,196
Digoxin	24 (3,6)	20 (2,9)	0,438

ACE inh: Angiotensin-converting enzyme inhibitor, ARB: Angiotensin II receptor blocker, NTG: Nitroglycerin